


Claims

1. A serine protease inhibitor of formula (I):



(I)

wherein:

R₂ is a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, oxygen or sulphur ring atom, optionally being substituted in the 3 and/or 4 position (in relation to the point of attachment of X-X) by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, MeSO₂- or R₁, or the substituents at the 3 or 4 positions taken together form a fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring optionally substituted by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}, and optionally substituted in the position alpha to the X-X group (i.e. 6 position for a six membered aromatic ring etc) by amino, hydroxy, halo, alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio with the proviso that R₂ cannot be aminoisoquinolyl;

each X independently is a C, N, O or S atom or a CO, CR_{1a}, C(R_{1a})₂ or NR_{1a} group, at least one X being C, CO, CR_{1a} or C(R_{1a})₂;

each R_{1a} independently represents hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkylaminocarbonyl, alkoxycarbonylamino, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

R₁ is as defined for R_{1a}, provided that R₁ is not

unsubstituted aminoalkyl;

Y (the α -atom) is a nitrogen atom or a CR_{1b} group;

Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group, optionally substituted by groups

5 R_{3a} or $R_{3i}X_i$;

each R_{3a} independently is R_{1c} , amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkylimidazolyl, thiazolyl, alkylthiazolyl, alkylloxazolyl, oxazolyl, alkylsulphonamido, 10 alkylaminosulphonyl, aminosulphonyl, haloalkoxy, haloalkyl, a group of the formula $-C(X^3)N(R^{11})R^{12}$ (wherein X^3 is O or S; and R^{11} and R^{12} are independently selected from hydrogen, methyl or ethyl or together with the nitrogen atom to which they are attached form a pyrrolidin-1-yl, piperidin-1-yl or 15 morpholino group), or $-OCH_2O-$ which is bonded to two adjacent ring atoms in Cy;

X_i is a bond, O, NH or CH_2 ;


R_{3i} is phenyl, pyridyl or pyrimidinyl optionally substituted by R_{3a} ; and

20 R_{1b} , R_{1c} and R_{1j} are as defined for R_{1a} ;

L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group; and

$Lp(D)_n$ is of the formula:

25



in which:

r is 1 or 2;

X_a is CH and X_b is N;

30 s, t and u are each 0 or 1;


L_a and L_b are each independently selected from a single bond, $C=O$, O and NR_{1e} , in which R_{1e} is hydrogen or (1-

6C)alkyl;

G is (1-6C)alkanediyl; and

R₁₀ is (1-6C)alkyl; (3-6C)cycloalkyl [which is unsubstituted or substituted by (1-6C)alkyl]; indanyl;

5 pyridyl; tetrahydropyranyl; tetrahydrothiopyranyl; phenyl {which is unsubstituted or substituted by one or two R₃ groups [wherein R₃ is hydrogen, hydroxyl, alkoxy, alkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), hydroxyalkyl (optionally substituted by hydroxy, 10 alkylamino, alkoxy, oxo, aryl or cycloalkyl), alkoxyalkyl, alkoxycarbonyl, alkylaminocarbonyl, alkoxycarbonylamino, acyloxymethoxycarbonyl, aminoalkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), alkylamino (optionally substituted by hydroxy, alkylamino, 15 alkoxy, oxo, aryl or cycloalkyl), amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkyl imidazolyl, thiazolyl, alkyl thiazolyl, alkyl oxazolyl, oxazolyl, alkylsulphonamido, alkylaminosulphonyl, aminosulphonyl, 20 haloalkoxy, or haloalkyl]}, pyrrolinyl; or a group of formula:



in which v is 1,2 or 3; one of X_c and X_d is N and the other is CH or N (provided that when v is 1, X_c and X_d are not both N); 25 and R₁₁ is hydrogen, (1-6C)alkyl or when X_d is CH, hydroxy(1-6C)alkyl; provided that when t is 0, the sum of s and u is 1; when X_b is N, L_a is a bond or C=O; when X_c is N, L_b is a bond or C=O; when X_b and X_c are both N, t is 1; and when (L_a)_s-(G)_t-(L_b)_u represents an alkyl group and X_b and X_c both 30 represent N, the alkyl group contains at least two chain carbon atoms;


or R₁₀ is hydrogen and s, t and u are each 0;

or the compound of formula (I) that is 4-{[4-methoxybenzoyl-D,L-(2-trifluoromethylthiophenyl)-glycinyllaminomethyl}-1-isopropylpiperidine;

or a physiologically-tolerable salt thereof.

5

2. A serine protease inhibitor of formula (I):



10 wherein:

R₂ is a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, oxygen or sulphur ring atom, optionally being substituted in the 3 and/or 4 position (in relation to the point of attachment of X-X) by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, MeSO₂- or R₁, or the substituents at the 3 or 4 positions taken together form a fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring optionally substituted by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}, and optionally substituted in the position alpha to the X-X group (i.e. 6 position for a six membered aromatic ring etc) by amino, hydroxy, halo, alkyl, carboxy, alkoxy, alkoxy, alkoxy, alkoxy or alkylthio with the proviso that R₂ cannot be aminoisquinolyl;

each X independently is a C, N, O or S atom or a CO,
CR_{1a}, C(R_{1a})₂ or NR_{1a} group, at least one X being C, CO, CR_{1a}
or C(R_{1a})₂;

30 each R_{1a} independently represents hydrogen or hydroxyl,
alkoxy, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl,
alkoxycarbonyl, alkylaminocarbonyl, alkoxyalkylamino,

acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

R_1 is as defined for R_{1a} , provided that R_1 is not unsubstituted aminoalkyl;

5 Y (the α -atom) is a nitrogen atom or a CR_{1b} group;


Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group optionally substituted by groups R_{3a} or phenyl optionally substituted by R_{3a} ;

each R_{3a} independently is R_{1c} , amino, halo, cyano, nitro, 10 thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkyl imidazolyl, thiazolyl, alkyl thiazolyl, alkyl oxazolyl, oxazolyl, alkylsulphonamido, alkylaminosulphonyl, aminosulphonyl, haloalkoxy or haloalkyl; and

15 R_{1b} , R_{1c} and R_{1j} are as defined for R_{1a} ;

L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group; and

$Lp(D)_n$ is of the formula:



20 in which:

r is 1 or 2;


X_a is CH and X_b is N;

s , t and u are each 0 or 1;

25 L_a and L_b are each independently selected from a single bond, C=O, O and NR_{1e} , in which R_{1e} is hydrogen or (1-6C)alkyl;

G is (1-6C)alkanediyl; and

R₁₀ is (1-6C)alkyl; (3-6C)cycloalkyl [which is unsubstituted or substituted by (1-6C)alkyl]; indanyl; pyridyl; tetrahydropyranyl; tetrahydrothiopyranyl; phenyl {which is unsubstituted or substituted by one or two R₃ groups [wherein R₃ is hydrogen, hydroxyl, alkoxy, alkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), hydroxyalkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), alkoxyalkyl, alkoxycarbonyl, alkylaminocarbonyl, alkoxycarbonylamino, acyloxymethoxycarbonyl, aminoalkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), alkylamino (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkyl imidazolyl, thiazolyl, alkyl thiazolyl, alkyl oxazolyl, oxazolyl, alkylsulphonamido, alkylaminosulphonyl, aminosulphonyl, haloalkoxy or haloalkyl]}, pyrrolinyl; or a group of formula:



20 in which v is 1, 2 or 3; one of X_C and X_D is N and the other is CH or N, provided that when v is 1, X_C and X_D are not both N; and R_{11} is hydrogen, (1-6C)alkyl or when X_D is CH, hydroxy(1-6C)alkyl; provided that when t is 0, the sum of s and u is 1; when X_B is N, L_A is a bond or C=O; when X_C is N, 25 L_B is a bond or C=O; when X_B and X_C are both N, t is 1; and when $(L_A)_S-(G)_T-(L_B)_U$ represents an alkyl group and X_B and X_C both represent N, the alkyl group contains at least two chain carbon atoms.


or a physiologically-tolerable salt thereof.

30 3. A serine protease inhibitor according to claim 1 or claim

2, wherein R³ is selected from hydrogen, hydroxyl, methoxy, ethoxy, methyl, ethyl, propyl, 2-propyl, butyl, 2-butyl, t-butyl, pentyl, 2-pentyl, 3-pentyl, isopropylaminomethyl, dimethylamino-methyl, diethylaminomethyl, dimethylaminoethyl, 5 acetyl, hydroxymethyl, hydroxyethyl, carboxy, carboxy(1-5C)alkyl, methoxymethyl, methoxycarbonyl, ethoxycarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, aminomethyl, aminocarbonyl, aminocarbonyl(1-5C)alkyl, methylamino, dimethylamino, ethylamino, formylamino, acetylamino, amino, 10 fluoro, chloro, cyano, nitro, thiol, methylthio, methylsulphonyl, ethylsulphonyl, isopropylsulphonyl, methylsulphenyl, 1,2,4-triazol-2-yl, 1,2,4-triazol-4-yl, 1,2,3-triazol-4-yl, 1,3-imidazol-1-yl, 1,3-imidazol-4-yl, tetrazol-1-yl, tetrazol-5-yl, methylsulphonamido, ethylsulphonamido, 15 propylsulphonamido, methylaminosulphonyl, ethylaminosulphonyl, propylaminosulphonyl, aminosulphonyl, trifluoromethoxy, trifluoromethyl and trichloromethyl.

4. A compound according to any of claims 1 to 3 wherein r is 20 2.

5. A compound according to claim 1 wherein Ip(D)_n is of the formula:



25 wherein:

q is 1 or 2;


R_s is hydrogen, -(CH₂)_c-R_c, -CHR_eR_f, or -CH₂-CHR_eR_f [c is 0, 1 or 2; wherein R_c is pyridyl or phenyl (which phenyl may bear a fluoro, chloro, methyl, CONH₂, SO₂NH₂,

methylaminosulphonyl, dimethylaminosulphonyl, methylsulphonylamino, methoxy or methylsulphonyl substituent) and R_e and R_f are independently hydrogen or C_{1-3} alkyl; or CHR_eR_f is (3-6C)cycloalkyl (which may bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position, provided the substituent is not bonded to the CH group which is bonded to L), tetrahydropyranyl, tetrahydrothiopyranyl, pyrrolidinyl (which may bear a 1-methyl substituent), piperidinyl (which may bear a 1-methyl substituent) (provided that the tetrahydropyranyl, tetrahydrothiopyranyl, pyrrolidinyl and piperidinyl rings are not linked to the piperidin-1,4-diyl group through a ring nitrogen atom or a ring carbon atom adjacent to a ring oxygen, sulfur or nitrogen atom) or indan-2-yl.

15

6. A compound according to any one of claims 1 to 5 wherein L is CONH, CH_2NHCO , $CONHCH_2$, $CONHCH_2CH_2$ or $CON(Me)CH_2$.

7. A serine protease inhibitor according to claim 2 wherein -L-Lp(D)_n is of the formula:



20

wherein

q is 1 or 2;


s is 0 or 1; and


R_s is $-(CH_2)_c-R_c$, $-CHR_eR_f$, or $-CH_2-CHR_eR_f$ [wherein c is 1 or 2; R_c is pyridyl or phenyl (which phenyl may bear a fluoro, chloro, methyl, $CONH_2$, SO_2NH_2 , methylaminosulphonyl, dimethylaminosulphonyl, methylsulphonylamino, methoxy or methylsulphonyl substituent) and R_e and R_f are independently

hydrogen or C_{1-3} alkyl; or CHR_eR_f is cyclopentyl (which may bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position), cyclohexyl (which may bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position),
5 tetrahydropyran-4-yl, tetrahydrothiopyran-4-yl, pyrrolidin-3-yl (which may bear a 1-methyl substituent), piperidin-4-yl (which may bear a 1-methyl substituent), or indan-2-yl].

8. A compound according to any of claims 5 to 7 wherein q is
10 2.

9. A compound according to claim 1 or claim 2 wherein $Lp(D)_n$ is selected from one of the following formulae:






wherein m represents 0 or 1.

- 5 10. A compound according to any of claims 5 to 7 wherein R_8
 is selected from: hydrogen, methyl, ethyl, prop-2-yl, but-2-
 yl, pent-3-yl, hept-4-yl, cyclopentyl, cyclohexyl,
 cyclohexylmethyl, 1-methylpiperidin-4-yl, tetrahydropyran-4-
 yl, tetrahydrothiopyran-4-yl, phenyl, benzyl, pyrid-2-yl,
 10 pyrid-3-yl, pyrid-4-yl, pyrid-3-ylmethyl, pyrid-4-ylmethyl and
 indan-2-yl.

11. A compound according to any one of claims 1 to 10 wherein
 R_2 is phenyl, thien-2-yl, naphthyl, indol-2-yl, indol-6-yl,
 15 benzo[b]furan-5-yl, benzo[b]thiophen-2-yl or benzimidazol-2-yl
 (each of which is optionally substituted as defined in claim
 1).

12. A compound according to any one of claims 1 to 11 wherein
 20 optional substituents for R_2 are selected from:
 fluoro, chloro, bromo, iodo, nitro, thiol, difluoromethoxy,
 trifluoromethoxy, hydrazido, methylhydrazido, amino, cyano,
 trifluoromethyl, methylthio, vinyl, ethynyl, acetylamino,
 carboxy, acetoxy, hydroxy, methyl, ethyl, amido ($CONH_2$),
 25 aminomethyl, methoxy and ethoxy.

13. A compound according to any one of claims 1 to 12 wherein
 R_2 is selected from one of the formula (A') to (H'):



wherein X_4 is O or S, R_{13} is selected from hydrogen, fluoro, chloro or methyl and R_{14} is selected from hydrogen, methyl, ethyl, fluoro, chloro, and methoxy and R_{15} is selected from hydrogen, methyl, fluoro, chloro and amino.

14. A compound according to claims 1 to 13, wherein R_2 is 4-methoxyphenyl, 3-amino-4-chlorophenyl, indol-2-yl, 5-chloroindol-2-yl, indol-6-yl, 3-chloroindol-6-yl or 3-methylindol-6-yl.

15. A compound according to any one of claims 1 to 14 wherein -X-X- is -CONH-.

16. A compound according to any one of claims 1 to 15 wherein Y is CH.
17. A compound according to any one of claims 1 to 16 wherein Cy is an optionally R_{3a} substituted: phenyl, pyridyl, thienyl, thiazolyl, naphthyl, piperidinyl, furanyl, pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, oxazolyl, imidazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, pyrimidinyl, 10 pridaziny, quinolyl, isoquinolyl, benzofuryl, benzothienyl or cycloalkyl group, or a phenyl group substituted by R_{3i}X_i in which X_i is a bond, O, NH or CH₂ and R_{3i} is phenyl, pyridyl or pyrimidinyl optionally substituted by R_{3a}.
18. A compound according to any one of claims 1 to 17 wherein Cy is an optionally R_{3a} substituted: phenyl, pyridyl, thienyl, thiazolyl, naphthyl, piperidinyl or cycloalkyl group.
19. A compound according to any one of claims 1 to 18 wherein 20 R_{3a} is selected from hydrogen, hydroxyl, alkoxy, alkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), aminoalkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), hydroxyalkyl (optionally substituted by hydroxy, alkylamino, 25 alkoxy, oxo, aryl or cycloalkyl), alkoxyalkyl, alkoxycarbonyl, alkylaminocarbonyl, alkoxycarbonylamino, alkylamino (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), for amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, alkylsulphonamido, 30 alkylaminosulphonyl, aminosulphonyl, haloalkoxy, haloalkyl, a group of the formula -C(X³)N(R¹¹)R¹² (wherein X³ is O or S; and R¹¹ and R¹² are independently selected from hydrogen, methyl or ethyl or together with the nitrogen atom to which they are attached form a pyrrolidin-1-yl, piperidin-1-yl or

morpholino group) and $-OCH_2O-$ which is bonded to two adjacent ring atoms in Cy.

20. A compound according to any one of claims 1 to 19 wherein
- 5 R_{3a} is selected from hydrogen, hydroxyl, alkoxy, alkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), hydroxyalkyl (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), alkoxyalkyl, alkoxycarbonyl, alkylaminocarbonyl,
- 10 alkoxycarbonylamino, alkylamino (optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), aminoalkyl (substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl), halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, alkylsulphonamido,
- 15 alkylaminosulphonyl, aminosulphonyl, haloalkoxy and haloalkyl.

21. A compound according to any one of claims 1 to 19 wherein R_{3a} is selected from hydrogen, hydroxyl, methoxy, ethoxy, methyl, ethyl, methylaminomethyl, dimethylaminomethyl,
- 20 hydroxymethyl, carboxy, methoxymethyl, methoxycarbonyl, ethoxycarbonyl, methylaminocarbonyl, dimethylamino-carbonyl, aminomethyl, $CONH_2$, CH_2CONH_2 , acetylamino, methoxycarbonylamino, ethoxycarbonylamino, t-butoxycarbonylamino, amino, fluoro, chloro, bromo, cyano,
- 25 nitro, thiol, methylthio, methylsulphonyl, ethylsulphonyl, methylsulphenyl, methylsulphonylamido, ethylsulphonylamido, methylaminosulphonyl, ethylaminosulphonyl, aminosulphonyl, trifluoromethoxy, trifluoromethyl, pyrrolidin-1-ylcarbonyl, piperidin-1-ylcarbonyl, morpholin-1-ylcarbonyl and $-OCH_2O-$
- 30 (which is bonded to two adjacent ring atoms in Cy).


22. A compound according to any one of claims 1 to 19 wherein R_{3a} is selected from hydrogen, hydroxyl, methoxy, ethoxy, methyl, ethyl, methylaminomethyl, dimethylaminomethyl,

hydroxymethyl, carboxy, methoxymethyl, methoxycarbonyl, ethoxycarbonyl, methylaminocarbonyl, dimethylamino-carbonyl, aminomethyl, CONH_2 , CH_2CONH_2 , acetyl amino, methoxycarbonylamino, ethoxycarbonylamino, t-

- 5 butoxycarbonylamino, amino, fluoro, chloro, cyano, nitro, thiol, methylthio, methylsulphonyl, ethylsulphonyl, methylsulphenyl, methylsulphonylamido, ethylsulphonylamido, methylaminosulphonyl, ethylaminosulphonyl, aminosulphonyl, trifluoromethoxy and trifluoromethyl.

10

23. A compound according to any one of claims 1 to 22 wherein Cy is selected from:



15

wherein:

X' is selected from O, S and NMe;

X'' is selected from O and S;

X''' is selected from O, S, NH and NMe;

Y' is selected from hydrogen, amino and methyl;

R_O is selected from hydrogen, methyl, fluoro, chloro, trifluoromethyl, methoxy, methylthio, methylsulphinyl and
5 methylsulphonyl;

R_m is selected from hydrogen, methyl, fluoro, chloro, trifluoromethyl, methoxy, methylthio, methylsulphinyl, methylsulphonyl, carboxy, methoxycarbonyl and a group of the formula -C(X³)N(R¹¹)R¹² (wherein X³ is O or S and R¹¹ and R¹²
10 are independently selected from hydrogen, methyl or ethyl or together with the nitrogen atom to which they are attached form a pyrrolidin-1-yl, piperidin-1-yl or morpholino group);
R_p is selected from hydrogen and fluoro; or
R_O and R_m or R_m and R_p form an -OCH₂O- group; or
15 R_O and R_m together with the ring to which they are attached form a 5 or 6 membered aryl or heteroaryl ring (wherein the heteroaryl ring contains 1 or 2 heteroatoms selected from nitrogen, oxygen and sulfur);

one of R_{O1} and R_{O2} is hydrogen and the other is R_O;

20

24. A compound according to any one of claims 1 to 19 wherein Cy is selected from phenyl, 2-chlorophenyl, 2-methoxyphenyl, 4-carbamoylphenyl, pyrid-2-yl, pyrid-3-yl, thien-2-yl, thien-3-yl, furan-2-yl, furan-3-yl, imidazol-2-yl, thiazol-2-yl,
25 thiazol-4-yl, thiazol-5-yl, naphthyl, isoquinolin-5-yl, isoquinolin-8-yl, quinolin-4-yl, quinolin-5-yl, and quinolin-8-yl.

25. A compound as claimed in any one of Claims 1 to 24, in
30 which the alpha atom in Y is carbon and has the conformation that would result from construction from a D-α-aminoacid NH₂-CR_{1b}(Cy)-COOH where the NH₂ represents part of X-X

26. A pharmaceutical composition, which comprises a compound

[illegible][illegible][illegible][illegible][illegible][illegible]

of the same kind as the one which has been found in the case of the other two.

[illegible]